

United States Patent [19]

Haga et al.

US005229403A

[11] Patent Number: 5,229,403

[45] Date of Patent: Jul. 20, 1993

[54] DIAMINOTRIFLUOROMETHYLPYRIDINE DERIVATIVES AND PHOSPHOLIPASE A₂ INHIBITOR CONTAINING THEM

[75] Inventors: Takahiro Haga; Hideo Sugi; Itaru Shigehara; Shinji Odawara; Syuichi Yotsuya; Hirohiko Kimura; Kazuhiro Yamamoto, all of Kusatsu, Japan

[73] Assignee: Ishihara Sangyo Kaisha Ltd., Osaka, Japan

[21] Appl. No.: 723,377

[22] Filed: Jun. 28, 1991

[30] Foreign Application Priority Data

Jul. 10, 1990 [JP] Japan 2-181999

[51] Int. Cl.⁵ A61K 31/44; C07D 213/75

[52] U.S. Cl. 514/352; 514/353; 544/131; 544/316; 544/355; 544/405; 546/112; 546/146; 546/172; 546/192; 546/193; 546/194; 546/261; 546/262; 546/268; 546/285; 546/305; 546/306; 546/307; 546/308; 546/311

[58] Field of Search 546/285, 308, 268, 305, 546/306; 514/352, 353

[56] References Cited

U.S. PATENT DOCUMENTS

3,746,531	7/1973	Doherty	71/94
3,961,063	6/1976	Parish	424/263
3,962,263	6/1976	Doherty	71/94
3,963,660	6/1976	Hall et al.	546/305
4,000,285	12/1976	Parish	424/263
4,002,761	1/1977	Parish	424/263

FOREIGN PATENT DOCUMENTS

1445677 12/1968 Fed. Rep. of Germany
2205194 8/1972 Fed. Rep. of Germany

OTHER PUBLICATIONS

Chemical Abstracts, vol. 100, No. 19, May 7, 1984, p. 526, 156508R, & JP-A-58-198469, Nov. 18, 1983, "Pyridyl-Hydrazide Compounds".


Primary Examiner—C. Warren Ivy

Assistant Examiner—James H. Turnipseed

Attorney, Agent, or Firm—Oblon, Spivak, McClelland, Maier & Neustadt

[57] ABSTRACT

A diaminotrifluoromethylpyridine derivative of the formula (I) or its salt:



(I)

wherein X is $-CW^1R^1$, $-COCOR^2$, $-CW^1NHCOR^2$, $-C(=W^1(W^2W^3$ or $-CW^1N(R^4)R^5$, and Y is alkyl, $-CW^3R^6$, $-COCOR^7$, $-NHCOR^7$, $-C(=W^3)W^4R^8$, $-(NH)_mSO_2R^9$, $-(NH)_mSO_2OR^{10}$ or $-(NH)_mSO_2N R^{11}R^{12}$, wherein each of R^1 , R^6 and R^9 , which are independent from one another, is a chain hydrocarbon group which may be substituted, a monocyclic hydrocarbon group which may be substituted, a polycyclic hydrocarbon group which may be substituted, a monocyclic heterocycle group which may be substituted or a polycyclic heterocycle group which may be substituted, each of R^2 and R^7 , which are independent from each other, is alkyl which may be substituted, alkoxy which may be substituted, phenyl which may be substituted or phenoxy which may be substituted, each of R^3 , R^8 and R^{10} , which are independent from one another, is alkyl which may be substituted, alkenyl which may be substituted, alkynyl which may be substituted, cycloalkyl which may be substituted, phenyl which may be substituted or benzyl which may be substituted, each of R^4 , R^5 , R^{11} and R^{12} , which are independent from one another, is alkyl which may be substituted, each of W^1 , W^2 , W^3 and W^4 , which are independent from one another, is an oxygen atom or a sulfur atom, and m is 0 or 1, provided that a combination wherein one of X and Y is $-COCF_2X^1$ wherein X^1 is a hydrogen atom, a halogen atom, alkyl or haloalkyl, and the other is $-COCF_2X^2$ wherein X^2 is a hydrogen atom, a halogen atom, alkyl, haloalkyl or alkylcarbonyl, or $-COOX^3$ wherein X^3 is alkyl which may be substituted or phenyl which may be substituted, is excluded.


5 Claims, No Drawings

DIAMINOTRIFLUOROMETHYLPYRIDINE DERIVATIVES AND PHOSPHOLIPASE A₂ INHIBITOR CONTAINING THEM

The present invention relates to novel diaminotrifluoromethylpyridine derivatives or their salts, a process for their production, a phospholipase A₂ inhibitor, an anti-inflammatory agent and an anti-pancreatitis agent containing them, and novel trifluoromethylpyridine derivatives as intermediates.

As a diaminotrifluoromethylpyridine derivative, for example, U.S. Pat. Nos. 3,746,531 and 3,962,263 disclose a pyridine as an active ingredient of a herbicide, which has trifluoromethyl at the 5-position, —NHCO—CF₂—T¹ wherein T¹ is a hydrogen atom, a chlorine atom, a fluorine atom, alkyl or haloalkyl at either the 2-position or the 3-position, and —NHCO—CF₂—T² wherein T² is a hydrogen atom, a chlorine atom, a fluorine atom, alkyl, haloalkyl or alkylcarbonyl, or —NHCOOT³ wherein T³ is C₁₋₄ lower alkyl or phenyl at the other position. However, this is different in the chemical structure from the diaminotrifluoromethylpyridine derivative of the present invention. Further, U.S. Pat. No. 3,961,063 discloses a trifluoromethyl-substituted pyridine as an active ingredient of an anthelmintic, which has —NHCSNHCO—T⁴ wherein T⁴ is alkoxy, at the 2- and 3-positions. However, this compound is different in the chemical structure from the diaminotrifluoromethylpyridine derivative of the present invention.

The present invention provides a diaminotrifluoromethylpyridine derivative of the formula (I) or its salt:



wherein X is —CW¹R¹, —COCOR², —CW¹NHCOR², —C(=W¹)W²R³ or —CW¹N(R⁴)R⁵, and Y is alkyl, —CW³R⁶, —COCOR⁷, —NHCOR⁷, —C(=W³)W⁴R⁸, —(NH)_mSO₂R⁹, —(NH)_mSO₂OR¹⁰ or —(NH)_mSO₂N(R¹¹)R¹², wherein each of R¹, R⁶ and R⁹, which are independent from one another, is a chain hydrocarbon group which may be substituted, a monocyclic hydrocarbon group which may be substituted, a polycyclic hydrocarbon group which may be substituted, a monocyclic heterocycle group which may be substituted or a polycyclic heterocycle group which may be substituted, each of R² and R⁷, which are independent from each other, is alkyl which may be substituted, alkoxy which may be substituted, phenyl which may be substituted or phenoxy which may be substituted, each of R³, R⁸ and R¹⁰, which are independent from one another, is alkyl which may be substituted, alkenyl which may be substituted, alkynyl which may be substituted, cycloalkyl which may be substituted, phenyl which may be substituted or benzyl which may be substituted, each of R⁴, R⁵, R¹¹ and R¹², which are independent from one another, is alkyl which may be substituted, each of W¹, W², W³ and W⁴, which are independent from one another, is an oxygen atom or a sulfur atom, and m is 0 or 1, provided that a combination wherein one of X and Y is —COCF₂X¹ wherein X¹ is a hydrogen atom, a halogen atom, alkyl or haloalkyl, and the other is —COCF₂X² wherein X² is a hydrogen atom, a halogen atom, alkyl, haloalkyl or alkylcarbonyl,

or —COOX³ wherein X³ is alkyl which may be substituted or phenyl which may be substituted, is excluded; a process for its production; a phospholipase A₂ inhibitor, an anti-inflammatory agent and an anti-pancreatitis agent containing it, and a trifluoromethylpyridine derivative as an intermediate.

Now, the present invention will be described in detail with reference to the preferred embodiments.

In the formula (I), the chain hydrocarbon group for each of R¹, R⁶ and R⁹ may be alkyl, alkenyl or alkynyl. The monocyclic hydrocarbon group may be cycloalkyl, cycloalkenyl or phenyl. The polycyclic hydrocarbon group may be a condensed polycyclic hydrocarbon group such as naphthyl, tetrahydronaphthyl or indanyl, or a bridged polycyclic hydrocarbon group such as adamantyl, noradamantyl, norbornanyl or norbornanonyl. The monocyclic heterocycle group may be pyrrolyl, furanyl, thienyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyrrolinyl, pyrrolidinyl, dihydrofuranyl, tetrahydrofuranyl, dihydrothienyl, tetrahydrothienyl, pyrazolinyl, hydantoinyl, oxazolinyl, isoxazolinyl, isoxazolidinyl, thiazolinyl, thiazolidinyl, dioxolanyl, dithiolanyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, dihydropyridyl, tetrahydropyridyl, piperidinyl, dihydrooxopyridazinyl, tetrahydrooxopyridazinyl, dihydrooxopyrimidinyl, tetrahydrooxopyrimidinyl, piperazinyl, dihydropyranyl, tetrahydropyranyl, dioxanyl, dihydrodithinyl, dithianyl or morphorinyl. The polycyclic heterocycle group may be a condensed polycyclic heterocycle group such as thienothienyl, dihydrocyclopentathienyl, indolyl, benzofuranyl, benzothienyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzimidazolyl, tetrahydrobenzothienyl, dihydrobenzofuranyl, tetrahydrobenzisoxazolyl, benzodioxolyl, quinolinyl, isoquinolinyl, benzodioxanyl or quinoxalinyl, or a bridged polycyclic heterocycle group such as quinuclidinyl.

The substituent for each of the chain hydrocarbon group which may be substituted for each of R¹, R⁶ and R⁹, the alkyl which may be substituted and the alkoxy which may be substituted for each of R² and R⁷, the alkyl which may be substituted, the alkenyl which may be substituted and the alkynyl which may be substituted for each of R³, R⁸ and R¹⁰, the alkyl which may be substituted for each of R⁴, R⁵, R¹¹ and R¹² and the alkyl which may be substituted for X³, may be a halogen atom, alkoxy, haloalkoxy, alkylthio, cycloalkyl, cycloalkoxy, cycloalkenyl, cycloalkenyloxy, alkoxy carbonyl, alkylcarbonyl, alkylcarbonyloxy, aryl, aryloxy, arylthio, amino or alkyl-substituted amino. The number of such substituents or substituents on such substituents may be one or more. When the number is two or more, such substituents may be the same or different.

The substituent for each of the monocyclic hydrocarbon group which may be substituted, the polycyclic hydrocarbon group which may be substituted, the monocyclic heterocycle group which may be substituted and the polycyclic heterocycle group which may be substituted for each of R¹, R⁶ and R⁹, the phenyl which may be substituted and the phenoxy which may be substituted for each of R² and R⁷, the cycloalkyl which may be substituted, the phenyl which may be substituted and the benzyl which may be substituted for each of R³, R⁸ and R¹⁰, and the phenyl which may be substituted for X³, may be a halogen atom, alkyl, haloalkyl, alkoxy, haloalkoxy, alkylthio, cycloalkyl, cycloalkoxy, cycloalkenyl, cycloalkenyloxy, alkoxy carbonyl,

alkylcarbonyl, alkylcarbonyloxy, aryl, aryloxy, arylthio, amino, alkyl-substituted amino, cyano or nitro. The number of such substituents or substituents for such substituents may be one or more. If the number is two or more, such substituents may be the same or different.

In the formula (I), the alkyl group and the alkyl moiety contained in each of X and Y may be C₁₋₁₈ alkyl such as methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, decyl or nonadecyl, and they include linear or branched aliphatic structural isomers. The alkenyl group and the alkenyl moiety contained in each of X and Y may be C₂₋₁₈ alkenyl such as vinyl, propenyl, butenyl, pentenyl, hexenyl, decenyl or nonadecenyl, and they include linear or branched aliphatic structural isomers. The alkynyl group and the alkynyl moiety contained in each of X and Y may be C₂₋₁₈ alkynyl such as ethynyl, propynyl, butynyl, pentynyl, hexynyl, decynyl or nonadecynyl, and they include linear or branched aliphatic structural isomers. The cycloalkyl group and the cycloalkyl moiety contained in each of X and Y may be C₃₋₈ cycloalkyl such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cyclooctyl. The cycloalkenyl group and the cycloalkenyl moiety contained in each of X and Y may be C₅₋₈ cycloalkenyl such as cyclopentenyl, cyclohexenyl or cyclooctenyl. The halogen atom contained in each of X and Y may be a fluorine atom, a chlorine atom, a bromine atom or an iodine atom. The aryl group and the aryl moiety contained in each of X and Y may be phenyl, thienyl, furanyl, pyridyl, naphthyl, benzothienyl, benzofuranyl or quinoliny.


Now, preferred embodiments of the compound of the present invention will be described. In the formula (I), it is preferred that X is —CW¹R¹ or —C(=W¹)W²R³, and Y is —SO₂R⁹. Each of R¹ and R⁶ is preferably alkyl which may be substituted, alkenyl which may be substituted, cycloalkyl which may be substituted, cycloalkenyl which may be substituted, phenyl which may be substituted, tetrahydronaphthyl which may be substituted, indanyl which may be substituted or thienyl which may be substituted, more preferably, alkyl, haloalkyl, alkenyl, haloalkenyl, cycloalkyl, halogen-substituted cycloalkyl, phenyl, halogen-substituted phenyl, alkyl- or haloalkyl-substituted phenyl, or alkoxy- or haloalkoxy-substituted phenyl. Each of R² and R⁷ is preferably alkoxy which may be substituted or phenyl which may be substituted, more preferably alkoxy, haloalkoxy, phenyl, or halogen-substituted phenyl. Each of R³, R⁸ and R¹⁰ is preferably alkyl which may be substituted, more preferably, alkyl or haloalkyl. Each of R⁴, R⁵, R¹¹ and R¹² is preferably alkyl. R⁹ is preferably alkyl which may be substituted, alkenyl which may be substituted, cycloalkyl which may be substituted, cycloalkenyl which may be substituted or phenyl which may be substituted, more preferably alkyl, haloalkyl, phenyl, halogen-substituted phenyl, alkyl- or haloalkyl-substituted phenyl, or alkoxy- or haloalkoxy-substituted phenyl.

Preferred specific compounds of the present invention include N-(2-ethylsulfonylamino-5-trifluoromethyl-3-pyridyl)cyclohexanecarboxamide, N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)-5-indanecarboxamide, N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)acetoxycetamide, N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)crotonamide, N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)-2-thiophenecarboxamide, N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)-3-trifluoromethylbenzamide, N-(2-ethylsulfonylamino-5-tri-


fluoromethyl-3-pyridyl)-3-fluorobenzamide, N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)-6-(1,2,3,4-tetrahydronaphthalene)carboxamide, N-(2-ethylsulfonylamino-5-trifluoromethyl-3-pyridyl)-crotonamide, N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)-3-(2-thienyl)acrylamide, and their salts.

The compound of the formula (I) may form a salt when Y is —SO₂R⁹ wherein R⁹ is as defined above. Such a salt may be any pharmaceutically acceptable salt, for example, an alkali metal salt such as a potassium salt or a sodium salt, an alkaline earth metal salt such as a calcium salt, or an organic amine salt such as a triethanol amine salt or a tris(hydroxymethyl)aminomethane salt. Such a salt may have crystal water.

The compounds of the formula (I) and (I-1) can be prepared, for example, by processes represented by the following reactions (A) and (B):



In the above formulas, R¹, R², R³, R⁴, R⁵, W¹, W², X and Y are as defined above, and Z is a halogen atom.



In the above formulas, Y¹ is —CW³R⁶, —COCOR⁷ or —C(=W³)W⁴R⁸, wherein R⁶, R⁷, R⁸, W³, W⁴, X and Z are as defined above.

A compound of the formula (I-1) wherein X and Y¹ are the same substituents, can be prepared in the same



5

manner as the Reaction (B) using as the starting material 2,3-diamino-5-trifluoromethylpyridine instead of the compound of the formula (III).

The reactions (A) and (B) are usually conducted in the presence of a solvent, if necessary, by using a base. The solvent may be an aromatic hydrocarbon such as benzene, toluene, xylene or chlorobenzene; a cyclic or non-cyclic aliphatic hydrocarbon such as chloroform, carbon tetrachloride, methylene chloride, dichloroethane, trichloroethane, n-hexane or cyclohexane; an ether such as diethyl ether, dioxane or tetrahydrofuran; a ketone such as acetone, methyl ethyl ketone or methyl isobutyl ketone; a nitrile such as acetonitrile or propionitrile; an aprotic polar solvent such as dimethylformamide, N-methylpyrrolidone, dimethylsulfoxide or sulfolane. The base may be an inorganic base or an organic base. The inorganic base may, for example, be an alkali metal hydroxide such as sodium hydroxide or potassium hydroxide; an alkali metal or alkaline earth metal carbonate such as anhydrous potassium carbonate or anhydrous calcium carbonate; an alkali metal hydride such as sodium hydride; or an alkali metal such as sodium metal. The organic base may be pyridine or triethylamine.


In the Reactions (A) and (B), a dehydrating condensation agent is required for the reaction with HOOCR^1 or HOOCR^6 . Such a dehydrating condensation agent may be dicyclohexylcarbodiimide, N,N'-carbonyldiimidazole or 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide. The reaction temperature is usually within a range of -30° to $+100^\circ$ C., preferably from 0° to 60° C., and the reaction time is usually within a range of from 1 to 24 hours, preferably from 1 to 10 hours.

The compound of the formula (II) can be prepared, for example, by processes represented by the following Reactions (C), (D) and (E):



6

-continued
Reaction (C)




In the above formulas, Y is as defined above.

The amination step in the above Reaction (C) is conducted usually in the presence of a solvent, if necessary, by using a base. The solvent may be an aromatic hydrocarbon such as benzene, toluene, xylene or chlorobenzene; a cyclic or non-cyclic aliphatic hydrocarbon such as chloroform, carbon tetrachloride, methylene chloride, dichloroethane, trichloroethane, n-hexane or cyclohexane; an ether such as diethyl ether, dioxane or tetrahydrofuran; a nitrile such as acetonitrile or propionitrile; or an aprotic polar solvent such as dimethylformamide, N-methylpyrrolidone, dimethylsulfoxide or sulfolane. The base may be the same as the one useful for the above-mentioned Reactions (A) and (B). The reaction temperature is usually within a range of from -30° to $+100^\circ$ C., and the reaction time is usually from 1 to 24 hours.

The reduction reaction in the reduction step in the above Reaction (C) may be conducted by a method wherein an acid such as hydrochloric acid or acetic acid is used together with iron or zinc, a method wherein sodium hydrosulfide, potassium hydrosulfide, sodium sulfide, potassium sulfide or sodium hydrosulfite is used, or a method of catalytic hydrogenation wherein hydrogen is used in the presence of a palladium catalyst or a nickel catalyst. The solvent to be used for the reduction may be optionally selected depending upon the reduction method. Usually, an alcohol such as methanol, ethanol or propanol, water, acetic acid, ethyl acetate, dioxane, tetrahydrofuran or acetonitrile may be employed. The reaction temperature is usually from 0° to 100° C., and the reaction time is usually from 1 to 24 hours.

(i) In a case where Y is $-\text{CW}^3\text{R}^6$ or $-\text{COCOR}^7$

-continued
Reaction (D)



In the above formulas, Y^2 is $-CW^3R^6$ or $-COR^7$, wherein R^6 , R^7 , W^3 and Z are as defined above.

The protecting group addition step and the Y^2 -modification step in the above Reaction (D) can be conducted in the same manner as in the above Reactions (A) and (B). Further, the protecting group removal step in the above Reaction (D) can be conducted by catalytic hydrogenation by means of a palladium catalyst such as palladium carbon usually in the presence of a solvent and an acid or base. The solvent may be water; an alcohol such as methanol or ethanol; or an ether such as diethyl ether, dioxane or tetrahydrofuran. The acid may be hydrobromic acid or trifluoroacetic acid. The base may be lithium hydroxide, potassium hydroxide, sodium hydroxide, potassium carbonate or sodium carbonate. The reaction temperature is usually from 0° to 100° C., and the reaction time is usually from 1 to 24 hours.

(ii) In a case where Y is $-SO_2R^9$


ganic base, for example, an alkali metal hydroxide such as sodium hydroxide or potassium hydroxide, or an alkali metal carbonate such as anhydrous potassium carbonate or anhydrous sodium carbonate. The reaction temperature is usually from 80° to 150° C., and the reaction time is usually from 1 to 10 hours.

The sulfonyl-modification step in the above Reaction (E) can be conducted in the same manner as in the above Reactions (A) and (B).

The nitration step in the above Reaction (E) can be conducted by reacting with nitric acid or nitrate usually in the presence of a solvent. The nitrate may be sodium nitrate or potassium nitrate. The solvent may be acetic acid, acetic anhydride or trifluoroacetic acid. The reaction temperature is usually from 50° to 120° C., and the reaction time is usually from 1 to 10 hours.

The reduction step in the above Reaction (E) can be conducted in the same manner as the reduction step in the above Reaction (C).




The compound of the above formula (III) can be



In the above formulas, Y^3 is $-SO_2R^9$, R^9 is alkyl which may be substituted, alkenyl which may be substituted, cycloalkyl which may be substituted or cycloalkenyl which may be substituted.

The amination step in the above Reaction (E) can be conducted usually in the presence of a solvent by means of a base. The solvent may be an aprotic polar solvent such as dimethyl acetamide, 1,3-dimethyl-2-imidazolidinone or dimethylsulfoxide. The base may be an inor-



prepared, for example, by a process represented by the following Reaction (F).



In the above formulas, $\text{R}^1, \text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{W}^1, \text{W}^2, \text{X}$ and Z are as defined above.

The above Reaction (F) can be conducted in the same manner as the above Reactions (A) and (B).





Among the compounds of the formula (IV), those wherein Y is $-\text{SO}_2\text{R}^9$, $-\text{SO}_2\text{OR}^{10}$ or $-\text{SO}_2\text{N}(\text{R}^{11})\text{R}^{12}$, can be produced also by a process represented by the following Reaction (G).



In the above formulas, Y^4 is $-\text{SO}_2\text{R}^9$, $-\text{SO}_2\text{OR}^{10}$ or $-\text{SO}_2\text{N}(\text{R}^{11})\text{R}^{12}$, wherein $\text{R}^9, \text{R}^{10}, \text{R}^{11}$ and R^{12} are as defined above.

The above Reaction (G) can be conducted in the same manner as the sulfonyl-modification step in the above Reaction (E).

The compound of the formula (I) can also be prepared by the following alternative method represented by a Reaction (H).




In the above formulas, $\text{R}^1, \text{R}^2, \text{R}^3, \text{R}^4, \text{R}^5, \text{W}^1, \text{W}^2, \text{X}$, Y and Z are as defined above.

The X-modification step in the above Reaction (H) can be conducted in the same manner as the above Reaction (A), and the amination step is conducted in the same manner as the amination step in the above Reaction (C).

Among the compounds of the above formulas (II), (IV), (IV-1), (V), (VI) and (VII), the following compounds are novel compounds and can be produced by the above Reactions (C), (E) and (G).

Trifluoromethylpyridine derivatives of the formula (VIII):



wherein Q is a hydrogen atom, nitro or amino, and Y^5 is $-(\text{NH})_m-\text{SO}_2\text{R}^9$ wherein R^9 and m are as defined above, $-(\text{NH})_m-\text{SO}_2\text{OR}^{10}$ wherein R^{10} and m are as defined above, or $-(\text{NH})_m-\text{SO}_2\text{N}(\text{R}^{11})\text{R}^{12}$ wherein $\text{R}^{11}, \text{R}^{12}$ and m are as defined above, provided that when Q is a hydrogen atom and m is 0, R^9 is other than naphthyl or phenyl which may be substituted.

Now, Preparation Examples for the compounds of the present invention will be described.

PREPARATION EXAMPLE 1

Preparation of

N-(2-ethylsulfonylamino-5-trifluoromethyl-3-pyridyl)-pentafluoropropionamide (Compound No. 19)

(1) 3.1 g of ethanesulfonamide was dissolved in 50 ml of dry tetrahydrofuran, and 1.2 g of 60% sodium hydride was added thereto under cooling with ice. After completion of the addition, the mixture was reacted for one hour under reflux. After cooling, 5.0 g of 2-chloro-3-nitro-5-trifluoromethylpyridine was added thereto, and then the mixture was reacted for 7 hours under reflux. After completion of the reaction, the reaction product was poured into 200 ml of water. Undissolved materials in water were extracted with ethyl ether and removed. Then, the aqueous layer was weakly acidified with dilute hydrochloric acid. Precipitated crystals were collected by filtration and dried to obtain 3.6 g of N-(3-nitro-5-trifluoromethyl-2-pyridyl)ethanesulfonamide having a melting point of from 160° to 163° C.

(2) 1.5 g of N-(3-nitro-5-trifluoromethyl-2-pyridyl)ethanesulfonamide obtained in the above step (1) was dissolved in 30 ml of methanol, and 0.2 g of 5% palladium/carbon was added thereto, and a reduction reaction

was conducted under a hydrogen pressure overnight under stirring. After completion of the reaction, 5% palladium/carbon was separated by filtration, and the solvent was distilled off under reduced pressure. The obtained crystals were washed with n-hexane and dried to obtain 1.2 g of N-(3-amino-5-trifluoromethyl-2-pyridyl)ethanesulfonamide having a melting point of from 118° to 120° C.

(3) 0.50 g of N-(3-amino-5-trifluoromethyl-2-pyridyl)ethanesulfonamide obtained in the above step (2) was suspended in 10 ml of dry diethyl ether, and 1.15 g of perfluoropropionic anhydride was dropwise added under cooling with ice. After the dropwise addition, the mixture was stirred for one hour and further reacted at room temperature for one hour. After completion of the reaction, the reaction product was poured into ice water and extracted with ethyl acetate. The extract layer was washed with water and dried, and the solvent was distilled off under reduced pressure. The obtained crystals were washed with n-hexane/ethyl ether to obtain 0.58 g of the desired product (Compound No. 19) having a melting point of from 168° to 170° C.

PREPARATION EXAMPLE 2

Preparation of

N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)-4-fluorobenzamide (Compound No. 10)

(1) 4.4 g of methanesulfonamide was dissolved in 70 ml of dry tetrahydrofuran, and 1.9 g of 60% sodium hydride was added thereto under cooling with ice. After completion of the addition, the mixture was reacted for one hour under reflux. After cooling, 7.0 g of 2-chloro-3-nitro-5-trifluoromethylpyridine was added thereto, and the mixture was reacted for 6 hours under reflux. After completion of the reaction, the reaction product was poured into 300 ml of water and washed with ethyl ether. Then, the aqueous layer was weakly acidified with dilute hydrochloric acid. Precipitated crystals were collected by filtration and dried to obtain 5.8 g of N-(3-nitro-5-trifluoromethyl-2-pyridyl)methanesulfonamide having a melting point of from 138° to 139° C.

(2) 4.0 g of N-(3-nitro-5-trifluoromethyl-2-pyridyl)methanesulfonamide obtained in the above step (1) was dissolved in 66 ml of methanol, and 0.4 g of 5% palladium/carbon was added thereto. A reduction reaction was conducted under a hydrogen pressure overnight under stirring. After completion of the reaction, 5% palladium/carbon was separated by filtration, and the solvent was distilled off under reduced pressure. The obtained crystals were washed with n-hexane and dried to obtain 3.2 g of N-(3-amino-5-trifluoromethyl-2-pyridyl)methanesulfonamide having a melting point of from 128° to 130° C.

(3) 0.50 g of N-(3-amino-5-trifluoromethyl-2-pyridyl)methanesulfonamide obtained in the above step (2) was dissolved in 6 ml of dry tetrahydrofuran, and 0.37 g of p-fluorobenzoyl chloride was dropwise added under cooling with ice. After the dropwise addition, the mixture was stirred for one hour and further reacted at room temperature overnight. After completion of the reaction, the reaction product was poured into ice water and extracted with ethyl acetate. The extract layer was washed with water and dried. The solvent was distilled off under reduced pressure, and the residue thereby obtained was crystallized from n-hexane/ethyl ether to obtain 0.61 g of the desired product (Com-

pound No. 10) having a melting point of from 211° to 213° C.

PREPARATION EXAMPLE 3

Preparation of

N-(3-trichloroacetyl-amino-5-trifluoromethyl-2-pyridyl)trifluoroacetamide (Compound No. 30)

(1) Into 38 ml of dry tetrahydrofuran, 1.5 g of 2,3-diamino-5-trifluoromethylpyridine was dissolved, and a solution mixture comprising 1.54 g of trichloroacetyl chloride and 3.8 ml of dry tetrahydrofuran was dropwise added thereto over a period of 10 minutes. Then, the mixture was reacted at room temperature for 3 hours. After completion of the reaction, precipitated crystals were collected by filtration and washed with tetrahydrofuran to obtain 2.2 g of N-(2-amino-5-trifluoromethyl-3-pyridyl)trichloroacetamide having a melting point of from 210° to 223° C.

(2) 2.20 g of N-(2-amino-5-trifluoromethyl-3-pyridyl)trichloroacetamide obtained in the above step (1) was dissolved in 45 ml of dry tetrahydrofuran, and a solvent mixture comprising 2.15 g of trifluoroacetic anhydride and 3 ml of dry tetrahydrofuran was dropwise added thereto under cooling with ice. After the dropwise addition, the mixture was reacted at room temperature for 3 hours. After completion of the reaction, the solvent was distilled off under reduced pressure, and the obtained crystals were washed with ethyl ether to obtain 1.20 g of the desired product (Compound No. 30) having a melting point of from 166° to 168° C.

PREPARATION EXAMPLE 4

Preparation of

N-(2-ethylsulfonylamino-5-trifluoromethyl-3-pyridyl)cyclohexanecarboxamide (Compound No. 47)

(1) 20.3 g of ethanesulfonamide and 26.0 g of 2-chloro-5-trifluoromethylpyridine were dissolved in 220 ml of dimethylsulfoxide, and 47.4 g of anhydrous potassium carbonate was further added thereto. This solution mixture was heated to 130° C. and reacted for 5 hours. After completion of the reaction, the reaction product was poured into 1 l of water. Undissolved materials in water were extracted with ethyl ether and removed. Then, the aqueous layer was adjusted to pH4 with concentrated hydrochloric acid, and precipitated crystals were collected by filtration and dried to obtain 26.2 g of N-(5-trifluoromethyl-2-pyridyl)ethanesulfonamide having a melting point of from 164° to 165° C.

(2) 45 g of N-(5-trifluoromethyl-2-pyridyl)ethanesulfonamide was dissolved in 112.5 ml of acetic acid. While heating it to a temperature of from 100° to 105° C., 26 g of fuming nitric acid (94%) was dropwise added, and the mixture was reacted for further 6 hours. The reaction product was left to cool to 80° C., and then poured into 2 l of ice water. Precipitated crystals were collected by filtration, washed with water and dried to obtain 47.8 g of N-(3-nitro-5-trifluoromethyl-2-pyridyl)ethanesulfonamide.

(3) 3.0 g of N-(3-nitro-5-trifluoromethyl-2-pyridyl)ethanesulfonamide was suspended in a solvent mixture comprising 30 ml of water and 30 ml of acetic acid, and 2.2 g of reduced iron was added thereto. Then, the mixture was heated to 50° C. and reacted for one hour. After completion of the reaction, the reaction product was cooled to room temperature, and excess iron was separated by filtration. The filtrate was extracted with

ethyl acetate. The extract layer was washed with water and dried. Ethyl acetate was distilled off under reduced pressure to obtain 2.5 g of N-(3-amino-5-trifluoromethyl-2-pyridyl)ethanesulfonamide.

An alternative process will be described. To a solution prepared by dissolving 34.9 g of sodium hydrosulfite in 400 ml of water, a solution prepared by dissolving 5.0 g of N-(3-nitro-5-trifluoromethyl-2-pyridyl)ethanesulfonamide in 80 ml of tetrahydrofuran, was dropwise added at room temperature. After completion of the dropwise addition, the mixture was reacted for further 3 hours. After completion of the reaction, sodium chloride was added until the tetrahydrofuran layer was separated. The separated tetrahydrofuran layer was dried, and tetrahydrofuran was distilled off under reduced pressure to obtain 4.2 g of N-(3-amino-5-trifluoromethyl-2-pyridyl)ethanesulfonamide.

(4) 2.36 g of N-(3-amino-5-trifluoromethyl-2-pyridyl)ethanesulfonamide was dissolved in 24 ml of dry tetrahydrofuran, and 1.54 g of cyclohexanecarbonyl chloride was dropwise added thereto under cooling with ice. After the dropwise addition, the mixture was stirred for one hour and further reacted at room temperature overnight. After completion of the reaction, the solvent was distilled off under reduced pressure, the obtained crystals were washed with ethyl ether to obtain 2.94 g of the desired product having a melting point of from 153° to 155° C.

An alternative process will be described. In 20 ml of methylene chloride, 0.5 g of 4-dimethylaminopyridine was dissolved, and 0.78 g of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride was added and dissolved. Then, 1 g of N-(3-amino-5-trifluoromethyl-2-pyridyl)ethanesulfonamide was added thereto, and 30 minutes later, 0.52 g of cyclohexanecarboxylic acid was added thereto, and stirring was conducted for 10 hours. After completion of the reaction, 40 ml of methylene chloride was added to the reaction product, and the reaction product was washed with 10% hydrochloric acid and then washed with an aqueous sodium chloride solution and then dried over anhydrous sodium sulfate. From the extract layer, solvent was distilled off and the obtained residue was purified by silica gel column chromatography to obtain 0.88 g of the desired product.

PREPARATION EXAMPLE 5

Preparation of sodium salt of N-(2-ethylsulfonylamino-5-trifluoromethyl-3-pyridyl)cyclohexanecarboxamide (Compound No. 251)






To 10 ml of an ethanol solution containing 1.00 g of N-(2-ethylsulfonylamino-5-trifluoromethyl-3-pyridyl)cyclohexanecarboxamide, 2.75 g of a 1N-sodium hydroxide aqueous solution was added under stirring at 40° C., and the mixture was stirred for one hour. After completion of the reaction, the solvent was distilled off under reduced pressure, and the obtained crystals were washed with ethyl ether to obtain 1.02 g of the desired product which decomposed at 299° C.

Trifluoromethylpyridine compounds of the above formula (VIII) are listed in Table 1.

TABLE 1













(VIII)

(VIII)

Intermediate No.	Q	Y ⁵	Melting point (°C.)
1	H	—SO ₂ CH ₃	189~191
2	H	—SO ₂ C ₂ H ₅	164~165
3	H	—SO ₂ CH ₂ CH ₂ CH ₃	157~159
4	H	—SO ₂ CH ₂ CH ₂ CH ₂ CH ₃	148~150
5	H	—SO ₂ CH(CH ₃) ₂	181~184
6	H	—SO ₂ CH(CH ₃)CH ₂ CH ₃	
7	H	—SO ₂ CH ₂ CH=CH ₂	
8	H	—SO ₂ CH ₂ CH ₂ CH(CH ₃) ₂	
9	H	—SO ₂ CH ₂ C(CH ₃)=CH ₂	
10	H	—SO ₂ CH ₂ CH ₂ OCH ₂ CH ₃	
11	H	—SO ₂ CF ₃	215~218
12	H	—SO ₂ — 	
13	H	—SO ₂ — 	
14	H	—SO ₂ — 	
15	H	—SO ₂ C ₈ H ₁₇ (n)	
16	H	—SO ₂ C ₁₈ H ₃₇ (n)	
17	H	—SO ₂ CF ₂ CF ₃	
18	NO ₂	—SO ₂ CH ₃	138~139
19	NO ₂	—SO ₂ CH ₂ CH ₃	160~163
20	NO ₂	—SO ₂ CH(CH ₃) ₂	138~140
21	NO ₂	—SO ₂ CH ₂ CH ₂ CH ₃	109~112
22	NO ₂	—SO ₂ CH ₂ CH ₂ CH ₂ CH ₃	76~78
23	NO ₂	—SO ₂ — 	138~140
24	NO ₂	—SO ₂ — 	145~146
25	NO ₂	—NH—SO ₂ CH ₃	175~182














15

TABLE 1-continued

(VIII)			
			
Intermediate No.	Q	Y ⁵	Melting point (°C.)
26	NO ₂		
27	NO ₂		
28	NO ₂		
29	NO ₂		51~56
30	NO ₂		156~158
31	NO ₂		
32	NO ₂		
33	NO ₂		
34	NO ₂		130~132
35	NO ₂		
36	NO ₂		

16

TABLE 1-continued













(VIII)			
			
Intermediate No.	Q	Y ⁵	Melting point (°C.)
37	NO ₂		192~194
38	NO ₂		
39	NO ₂		
40	NO ₂		
41	NO ₂		
42	NO ₂		
43	NO ₂		
44	NO ₂		
45	NO ₂		
46	NO ₂		
47	NO ₂		
48	NO ₂		148~149

17

TABLE 1-continued



















				(VIII)
Intermediate No.	Q	Y ⁵	Melting point (°C.)	
49	NO ₂		132	
50	NO ₂	-SO ₂ CF ₃	126 ~ 127	
51	NO ₂	-SO ₃ CH ₃	93 ~ 94	
52	NO ₂	-SO ₃ C ₂ H ₅	120 ~ 121	
53	NO ₂		104 ~ 105	
54	NO ₂			
55	NH ₂	-SO ₂ CH ₃	128 ~ 130	
56	NH ₂	-SO ₂ CH ₂ CH ₃	118 ~ 120	
57	NH ₂		155 ~ 157	
58	NH ₂	-SO ₂ CH ₂ CH ₂ CH ₃	82 ~ 84	
59	NH ₂	-SO ₂ CH ₂ CH ₂ CH ₂ CH ₃	102 ~ 103	
60	NH ₂		200 ~ 204	
61	NH ₂		170 ~ 175	
62	NH ₂	-NHSO ₂ CH ₃	128 ~ 133	
63	NH ₂			
64	NH ₂			
65	NH ₂			
66	NH ₂		136 ~ 139	

TABLE 1-continued

(VIII)			
			
Intermediate No.	Q	Y ⁵	Melting point (°C.)
78	NH ₂		
79	NH ₂		
80	NH ₂		
81	NH ₂		
82	NH ₂		
83	NH ₂		
84	NH ₂		
85	NH ₂		165~167
86	NH ₂		134~136
87	NH ₂	-SO ₂ CF ₃	122~124
88	NH ₂	-SO ₃ CH ₃	97~100
89	NH ₂	-SO ₃ C ₂ H ₅	131~132
90	NH ₂		223~227
91	NH ₂		



Compounds of the above formula (II) which are not included in the compounds of the above formula (VIII) are listed in Table 2.

TABLE 2

(II)			
			
Intermediate No.	Y	Melting point (°C.)	
5			
10			
15		207~210	
20	  	187~192 289~292	
25			
30			
35	   		
40			
45			
50			
55			
60			
65			
110			
111			
112			
113			
114			
115			
116			




21

TABLE 2-continued

<div style="display: flex; align-items: center; justify-content: center;"> <div style="text-align: center;">  </div> <div style="margin-left: 20px;">(II)</div> </div>		
Intermediate No.	Y	Melting point (°C.)
117		5




Compounds of the above formula (III) are listed in Table 3.

TABLE 3

<div style="display: flex; align-items: center; justify-content: center;"> <div style="text-align: center;">  </div> <div style="margin-left: 20px;">(III)</div> </div>		
Intermediate No.	X	Melting point (°C.)
118	-COCHCl ₂	170~171
119	-COCCl ₃	141~143
120	-COOCH ₂ CH ₃	151~154
121		156~158
122	-COCOCH ₃	
123		
124	-CONHCOCH ₃	



22

TABLE 3-continued

<div style="display: flex; align-items: center; justify-content: center;"> <div style="text-align: center;">  </div> <div style="margin-left: 20px;">(III)</div> </div>		
Intermediate No.	X	Melting point (°C.)
125		10
126		248~251

Compounds of the above formula (IV) which are not included in the compounds of the above formula (VIII) are listed in Table 4.

TABLE 4

<div style="display: flex; align-items: center; justify-content: center;"> <div style="text-align: center;">  </div> <div style="margin-left: 20px;">(IV)</div> </div>		
Intermediate No.	Y ²	Melting point (°C.)
127		189~195
128	-NHCOOCH ₂ CH ₃	97~99
129	-NHCOCH ₃	
130	-CH ₃	
131	-CH ₂ CH ₃	

Typical specific examples of the compound of the formula (I) of the present invention are listed in Table 5.

TABLE 5



<div style="display: flex; align-items: center; justify-content: center;"> <div style="text-align: center;">  </div> <div style="margin-left: 20px;">(I)</div> </div>				
Compound No.	X	Y	Type of salt	Melting point (°C.)
1	-CO(CH ₂) ₂ CH ₃	-SO ₂ CH ₃		113~114
2	-CO(CH ₂) ₃ CH ₃	-SO ₂ CH ₃		119~121
3	-CO(CH ₂) ₄ CH ₃	-SO ₂ CH ₃		119~122
4	-CO(CH ₂) ₇ CH ₃	-SO ₂ CH ₃		99~101
5	-CO(CH ₂) ₁₀ CH ₃	-SO ₂ CH ₃		94~97
6	-CO(CH ₂) ₁₄ CH ₃	-SO ₂ CH ₃		99~103
7	-COCH ₂ C(CH ₃) ₃	-SO ₂ CH ₃		150~151
8		-SO ₂ CH ₃		110~116
9	-COCH=CH ₂	-SO ₂ CH ₃		174~176

TABLE 5-continued












				(I)	
Compound No.	X	Y	Type of salt	Melting point (°C.)	
10		-SO ₂ CH ₃		211~213	
11	-COCF ₂ Cl	-SO ₂ CH ₃		199~201	
12	-COCF ₃	-SO ₂ CH ₃		154~157	
13	-COCF ₂ CF ₃	-SO ₂ CH ₃		186~189	
14	-COCF ₂ CF ₂ CF ₃	-SO ₂ CH ₃		170~173	
15	-COOC ₂ H ₅	-SO ₂ CH ₃		180~182	
16	-COO(CH ₂) ₂ CH ₃	-SO ₂ CH ₃		173~176	
17	-COO(CH ₂) ₃ CH ₃	-SO ₂ CH ₃		127~129	
18	-CSNHCOOC ₂ H ₅	-SO ₂ CH ₃		More than 300	
19	-COCF ₂ CF ₃	-SO ₂ C ₂ H ₅		168~170	
20	-COCF ₂ Cl	-SO ₂ C ₂ H ₅		171~174	
21	-CSNHCOOC ₂ H ₅	-SO ₂ C ₂ H ₅		More than 300	
22	-COCF ₂ CF ₃	-SO ₂ C ₃ H ₇ (n)		129~133	
23	-COCF ₂ CF ₃	-SO ₂ C ₈ H ₁₇ (n)		109~112	
24	-COCF ₃			160~163	
25	-CSNHCOOC ₂ H ₅			195~200	
26	-CO(CH ₂) ₂ OC ₂ H ₅	-CO(CH ₂) ₂ OC ₂ H ₅		75~76	
27	-COCF ₃	-COCHCl ₂		117~119	
28	-COCHCl ₂	-COCHCl ₂		158~159	
29	-COCHCl ₂	-COCF ₃		177~178	
30	-COCCl ₃	-COCF ₃		166~168	
31		-SO ₂ C ₂ H ₅		135~137	
32		-COCF ₂ CF ₃		228~230	
33		-SO ₂ C ₂ H ₅		130~134	
34		-SO ₂ CH ₃		218~222	
35		-SO ₂ CH ₃		219~224	
36		-SO ₂ C ₂ H ₅			












TABLE 5-continued



(I)


Compound No.	X	Y	Type of salt	Melting point (°C.)
53		$-\text{SO}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$		152 ~ 154
54	$-\text{CSNHCOOC}_2\text{H}_5$	$-\text{CH}_3$		191 ~ 193
55	$-\text{COCH}=\text{CHCH}_3$	$-\text{SO}_2\text{CH}_3$		158 ~ 161
56		$-\text{SO}_2\text{C}_2\text{H}_5$		234 ~ 237
57		$-\text{SO}_2\text{CH}_3$		210 ~ 214
58		$-\text{SO}_2\text{CH}_3$		220 ~ 222
59	$-\text{CO}-\text{CF}_2\text{CF}_2\text{H}$	$-\text{SO}_2\text{C}_2\text{H}_5$		
60		$-\text{SO}_2\text{CH}_3$		163 ~ 166
61		$-\text{SO}_2\text{CH}_3$		172 ~ 174
62		$-\text{SO}_2\text{CH}_3$		147 ~ 148
63	$-\text{COCH}_2\text{OCOCH}_3$	$-\text{SO}_2\text{CH}_3$		155 ~ 156
64		$-\text{SO}_2\text{CH}_3$		163 ~ 165
65	$-\text{COCH}(\text{C}_2\text{H}_5)(\text{CH}_2)_3\text{CH}_3$	$-\text{SO}_2\text{CH}_3$		141 ~ 144
66		$-\text{SO}_2\text{CH}_3$		128 ~ 130
67		$-\text{SO}_2\text{CH}_3$		126 ~ 130

TABLE 5-continued

Compound No.		X	Y	Type of salt	Melting point (°C.)
		<div style="text-align: center;">  </div>			
68			$-\text{SO}_2\text{CH}_3$		143~145
69			$-\text{SO}_2\text{CH}_3$		176~179
70		$-\text{COCH}=\text{C}(\text{CH}_3)_2$	$-\text{SO}_2\text{CH}_3$		187~188
71			$-\text{SO}_2\text{CH}_3$		215~218
72			$-\text{SO}_2\text{CH}_3$		227~229
73		$-\text{COCH}=\text{CHCH}=\text{CHCH}_3$	$-\text{SO}_2\text{CH}_3$		300
74		$-\text{CO}(\text{CH}_2)_2\text{CH}=\text{CH}_2$	$-\text{SO}_2\text{CH}_3$		91~93
75			$-\text{SO}_2\text{CH}_3$		209~210
76			$-\text{SO}_2\text{CH}_3$		245~249
77			$-\text{SO}_2\text{CH}_3$		229~231
78			$-\text{SO}_2\text{CH}_3$		187~189
79			$-\text{SO}_2\text{CH}_3$		198~201
80			$-\text{SO}_2\text{CH}_3$		230~233

(I)

TABLE 5-continued



(I)

Compound No.	X	Y	Type of salt	Melting point (°C.)
81		-SO ₂ CH ₃		211 ~ 215
82		-SO ₂ CH ₃		206 ~ 210
83		-SO ₂ CH ₃		207 ~ 210
84		-SO ₂ CH ₃		202 ~ 205
85		-SO ₂ CH ₃		227 ~ 231
86		-SO ₂ CH ₃		250 ~ 252
87		-SO ₂ CH ₃		194 ~ 197
88		-SO ₂ CH ₃		229 ~ 233
89	-COCCl ₂ CH ₃	-SO ₂ CH ₃		212 ~ 214
90		-SO ₂ CH ₃		231 ~ 234
91		-SO ₂ CF ₃		175 ~ 178
92		-SO ₂ CF ₃		209 ~ 210
93	-COCH=CHCH ₃	-SO ₂ C ₂ H ₅		158 ~ 160
94		-SO ₂ C ₂ H ₅		157 ~ 161

TABLE 5-continued

(I)

Compound No.	X	Y	Type of salt	Melting point (°C.)
95		$-\text{SO}_2\text{C}_2\text{H}_5$		147~148
96		$-\text{SO}_2\text{C}_2\text{H}_5$		163~165
97		$-\text{SO}_2\text{C}_2\text{H}_5$		163~166
98		$-\text{SO}_2\text{C}_2\text{H}_5$		204~208
99		$-\text{SO}_2\text{C}_2\text{H}_5$		215~218
100		$-\text{SO}_2\text{C}_2\text{H}_5$		233~237
101		$-\text{SO}_2\text{C}_2\text{H}_5$		208~209
102		$-\text{SO}_2\text{C}_2\text{H}_5$		188~190
103		$-\text{SO}_2\text{C}_3\text{H}_7(\text{iso})$		152~154
104		$-\text{SO}_2\text{C}_3\text{H}_7(\text{iso})$		216~217
105		$-\text{SO}_2\text{C}_3\text{H}_7(\text{iso})$		227~230
106	$-\text{COOC}_3\text{H}_7(\text{n})$	$-\text{SO}_2\text{C}_3\text{H}_7(\text{iso})$		161~163

TABLE 5-continued














				(I)	
Compound No.	X	Y	Type of salt	Melting point (°C.)	
107		$-\text{SO}_2\text{C}_4\text{H}_9(n)$		138 ~ 139	
108	$-\text{COCF}_2\text{Cl}$	$-\text{SO}_2\text{C}_4\text{H}_9(n)$		156	
109				202 ~ 205	
110	$-\text{CO}(\text{CH}_2)_4\text{CH}_3$	$-\text{SO}_2\text{N}(\text{CH}_3)_2$		97	
111		$-\text{SO}_2\text{N}(\text{CH}_3)_2$		168 ~ 169	
112	$-\text{COCF}_2\text{CF}_3$	$-\text{SO}_2\text{N}(\text{CH}_3)_2$		157 ~ 159	
113		$-\text{SO}_2\text{N}(\text{CH}_3)_2$		189 ~ 191	
114	$-\text{COOC}_3\text{H}_7(n)$	$-\text{SO}_2\text{N}(\text{CH}_3)_2$		174 ~ 176	
115		$-\text{SO}_2\text{OCH}_3$		147 ~ 148	
116		$-\text{SO}_2\text{OCH}_3$		163 ~ 164	
117		$-\text{SO}_2\text{OC}_2\text{H}_5$		140 ~ 141	
118		$-\text{SO}_2\text{OC}_2\text{H}_5$		160 ~ 162	
119		$-\text{SO}_2\text{C}_2\text{H}_5$		137 ~ 139	
120		$-\text{SO}_2\text{CH}_3$		202 ~ 203	
121		$-\text{SO}_2\text{CH}_3$		145 ~ 147	

TABLE 5-continued




















Compound No.		X		Y		Type of salt	Melting point (°C.)
							(I)
							
122				-SO ₂ CH ₃			221 ~ 224
123				-SO ₂ CH ₃			184 ~ 185
124		-CO(CH ₂) ₅ CH ₃		-SO ₂ CH ₃			94 ~ 96
125		-CO(CH ₂) ₆ CH ₃		-SO ₂ CH ₃			94 ~ 96
126							178 ~ 180
127							226 ~ 228
128				-SO ₂ CH ₃			
129				-SO ₂ CH ₃			
130				-SO ₂ CH ₃			
131				-SO ₂ CH ₃			
132				-SO ₂ CH ₃			
133							
134							
135							

TABLE 5-continued

(I)



Compound No.	X	Y	Type of salt	Melting point (°C.)
136		$-\text{SO}_2\text{C}_2\text{H}_5$		
137				
138				
139		$-\text{SO}_2\text{CH}_2-\text{C}(\text{CH}_3)=\text{CH}_2$		138 ~ 140
140		$-\text{SO}_2-\text{C}_6\text{H}_{11}$		190 ~ 192
141				
142		$-\text{SO}_2\text{C}_2\text{H}_5$		210 ~ 211
143		$-\text{SO}_2\text{C}_2\text{H}_5$		
144		$-\text{SO}_2\text{C}_2\text{H}_5$		
145				
146		$-\text{SO}_2\text{C}_2\text{H}_5$		

TABLE 5-continued



















					
Compound No.	X	Y	Type of salt	Melting point (°C.)	(I)
147		-SO ₂ C ₂ H ₅			
148		-SO ₂ C ₂ H ₅			
149		-SO ₂ C ₂ H ₅			
150					
151					
152		-SO ₂ C ₂ H ₅			
153					
154				166~167	
155				144~146	
156					
157		-SO ₂ C ₂ H ₅			

TABLE 5-continued

















Compound No.		X		Y		Type of salt	Melting point (°C.)
						(I)	
158				-SO ₂ C ₂ H ₅			
159							
160							
161				-SO ₂ C ₂ H ₅			
162							133~135
163				-SO ₂ C ₂ H ₅			
164				-SO ₂ C ₂ H ₅			
165				-SO ₂ C ₂ H ₅			
166				-SO ₂ C ₂ H ₅			
167							
168				-SO ₂ C ₂ H ₅			

TABLE 5-continued













					
				(I)	
Compound No.	X	Y	Type of salt	Melting point (°C.)	
169		-SO ₂ C ₂ H ₅			
170		-SO ₂ C ₂ H ₅			
171		-SO ₂ C ₂ H ₅			
172		-SO ₂ C ₂ H ₅			
173		-SO ₂ C ₂ H ₅			
174		-SO ₂ C ₂ H ₅			
175		-SO ₂ C ₂ H ₅			
176		-SO ₂ C ₂ H ₅			
177		-SO ₂ C ₂ H ₅			
178		-SO ₂ C ₂ H ₅			
179		-SO ₂ C ₂ H ₅			

TABLE 5-continued


























Compound No.		X		Y		Type of salt	Melting point (°C.)
							(I)
							
180				-SO ₂ C ₂ H ₅			
181				-SO ₂ C ₂ H ₅			
182				-SO ₂ C ₂ H ₅			
183				-SO ₂ C ₂ H ₅			
184							
185				-SO ₂ C ₂ H ₅			
186							
187				-SO ₂ C ₂ H ₅			
188				-SO ₂ C ₂ H ₅			
189				-SO ₂ C ₂ H ₅			

TABLE 5-continued

Compound No.		X	Y	Type of salt	Melting point (°C.)
<div style="text-align: center;">  </div>					
190			$-\text{SO}_2\text{C}_2\text{H}_5$		
191			$-\text{SO}_2-\text{N}$ 		
192			$-\text{SO}_2\text{C}_2\text{H}_5$		
193			$-\text{SO}_2\text{C}_2\text{H}_5$		
194			$-\text{SO}_2\text{C}_2\text{H}_5$		
195			$-\text{SO}_2\text{C}_2\text{H}_5$		
196			$-\text{SO}_2\text{C}_2\text{H}_5$		
197			$-\text{SO}_2\text{C}_2\text{H}_5$		
198			$-\text{SO}_2\text{C}_2\text{H}_5$		

(I)

TABLE 5-continued



(I)

Compound No.	X	Y	Type of salt	Melting point (°C.)
199		$-\text{SO}_2\text{C}_2\text{H}_5$		
200		$-\text{SO}_2\text{C}_2\text{H}_5$		
201		$-\text{SO}_2\text{C}_2\text{H}_5$		
202		$-\text{SO}_2\text{C}_2\text{H}_5$		
203		$-\text{SO}_2-\text{N}$		
204		$-\text{SO}_2\text{C}_2\text{H}_5$		
205		$-\text{SO}_2\text{C}_2\text{H}_5$		
206		$-\text{SO}_2\text{C}_2\text{H}_5$		
207		$-\text{SO}_2\text{C}_2\text{H}_5$		265 ~ 266
208				

TABLE 5-continued















Compound No.		X	Y	Type of salt	Melting point (°C.)
<div style="text-align: center;">  </div>					
209			$-\text{SO}_2\text{C}_2\text{H}_5$		
210			$-\text{SO}_2\text{C}_2\text{H}_5$		
211			$-\text{SO}_2\text{C}_2\text{H}_5$		
212					
213			$-\text{SO}_2\text{C}_2\text{H}_5$		
214			$-\text{SO}_2\text{C}_2\text{H}_5$		248 ~ 249
215			$-\text{SO}_2\text{C}_2\text{H}_5$		
216			$-\text{SO}_2\text{C}_2\text{H}_5$		
217			$-\text{SO}_2\text{C}_2\text{H}_5$		219 ~ 221
218			$-\text{SO}_2\text{C}_2\text{H}_5$		241 ~ 242
219					

TABLE 5-continued












				(I)	
Compound No.	X	Y	Type of salt	Melting point ("C.)	
220		$-\text{SO}_2\text{C}_2\text{H}_5$			
221					
222		$-\text{SO}_2\text{C}_2\text{H}_5$			
223		$-\text{SO}_2\text{C}_2\text{H}_5$			
224		$-\text{SO}_2\text{C}_2\text{H}_5$			
225	$-\text{C}(=\text{O})-\text{CH}_2-\text{OCH}_2\text{CF}_3$	$-\text{SO}_2\text{C}_2\text{H}_5$			
226	$-\text{C}(=\text{O})-\text{CH}_2\text{SCH}_3$	$-\text{SO}_2\text{C}_2\text{H}_5$			
227		$-\text{SO}_2\text{C}_2\text{H}_5$			
228		$-\text{SO}_2\text{C}_2\text{H}_5$			
229		$-\text{SO}_2\text{C}_2\text{H}_5$			
230	$-\text{C}(=\text{O})\text{CH}_2\text{COOCH}_3$	$-\text{SO}_2\text{C}_2\text{H}_5$			
231	$-\text{C}(=\text{O})(\text{CH}_2)_2\text{C}(=\text{O})\text{CH}_3$	$-\text{SO}_2\text{C}_2\text{H}_5$			
232		$-\text{SO}_2\text{C}_2\text{H}_5$			

TABLE 5-continued














Compound No.		X		Y		Type of salt	Melting point (°C.)
							(I)
							
233				-SO ₂ C ₂ H ₅			
234				-SO ₂ C ₂ H ₅			
235				-SO ₂ C ₂ H ₅			
236				-SO ₂ C ₂ H ₅			
237				-SO ₂ C ₂ H ₅			
238				-SO ₂ C ₂ H ₅			
239				-SO ₂ C ₂ H ₅			
240				-SO ₂ C ₂ H ₅			
241				-SO ₂ C ₂ H ₅			
242				-SO ₂ C ₂ H ₅			
243				-SO ₂ C ₂ H ₅			







TABLE 5-continued



(I)

Compound No.	X	Y	Type of salt	Melting point (°C.)
244		$-\text{SO}_2\text{C}_2\text{H}_5$		
245		$-\text{SO}_2\text{C}_2\text{H}_5$		
246		$-\text{SO}_2\text{CH}_2-$		
247		$-\text{SO}_2\text{C}_2\text{H}_5$		
248		$-\text{SO}_2-$		
249		$-\text{SO}_2\text{C}_3\text{H}_7(n)$		
250		$-\text{SO}_2\text{C}_3\text{H}_7(n)$		
251		$\text{SO}_2\text{C}_2\text{H}_5$	Na salt	299 (decomposed)
252		$\text{SO}_2\text{C}_2\text{H}_5$	K salt	More than 300
253		$\text{SO}_2\text{C}_3\text{H}_7(\text{iso})$	Na salt	
254		SO_2CF_3	Na salt	
255		SO_2-	Na salt	

TABLE 5-continued

Compound No.		X	Y	Type of salt	Melting point (°C.)
<div style="text-align: center;">  (I) </div>					
256			SO ₂ C ₂ H ₅	Na salt	
257			SO ₂ C ₂ H ₅	Na salt	
258			SO ₂ C ₂ H ₅	Na salt	
259			—SO ₂ CH ₃	Na salt	More than 300
260		—COCF ₂ CF ₃	—SO ₂ CH ₃	Na salt	More than 300
261		—COCF ₂ CF ₃	—SO ₂ C ₂ H ₅	Na salt	
262			—SO ₂ C ₂ H ₅	Ca salt	245 (decomposed)

The compound of the formula (I) of the present invention is useful as an active ingredient for a phospholipase A₂ inhibitor, an anti-inflammatory agent or an anti-pancreatitis agent. Phospholipase A₂ can be detected in various tissues or cells in a body. It is said that in platelets or cells related to inflammatory symptoms, phospholipase A₂ is secreted or activated by various stimulations and contributes to the production of a platelet activating factor (PAF) or some arachidonic acid metabolites. The arachidonic acid metabolites have been found to be closely related to various diseases, for example, inflammatory symptoms such as rheumatoid arthritis, arthritis deformans, tenonitis, psoriasis and related dermatitis; nasal and bronchial airway troubles such as allergic rhinitis and allergic bronchial asthma; and immediate hypersensitive reactions such as allergic conjunctivitis. On the other hand, phospholipase A₂ secreted from pancreas is activated in the intestine and exhibits a digestive action, but once activated in the pancreas, it is believed to be one of the factors causing pancreatitis. The compound of the present invention inhibits phospholipase A₂ and thus is effective for the treatment of the above-mentioned diseases caused by phospholipase A₂ such as inflammatory symptoms, nasal and bronchial airway troubles, immediate hypersensitive reactions or pancreatitis. Thus, it is useful as an anti-inflammatory agent, an agent for treating bronchial asthma, an anti-allergy agent, an anti-pancreatitis agent, anti-nephritis agent, or anti-MOFC (Multiple Organ Failure).

In regard to the efficacy against pancreatitis, the compound of the present invention is expected to be more efficient by using in combination with other drugs, for example, a proteinase inhibitor, such as galexate mesilate, camostat mesilate, or nafamostat mesilate.

The compound of the present invention is particularly suitable for use as an anti-inflammatory agent and/or an anti-pancreatitis agent.

TEST EXAMPLE 1

Phospholipase A₂ inhibitory activity, method A

(1) Preparation of substrate

To 10 mg of egg yolk lecithin (manufactured by Wako Pure Chemical Industries Ltd.), 1 ml of glycerine, 2 ml of a 50 mM Tris-HCl buffer solution (pH7.5) [Tris(hydroxymethyl)aminomethane (manufactured by Nacalai Tesque K.K.) was adjusted to pH7.5 with hydrochloric acid], 0.5 ml of a 150 mM calcium chloride solution (calcium chloride was dissolved in a 50 mM Tris-HCl buffer solution) and 0.5 ml of a 0.05% Triton-X100 (manufactured by Nacalai Tesque K.K.) solution (Triton-X100 was dissolved in a 50 mM Tris-HCl buffer solution), were added and dispersed by an agate mortar or dispersed by an ultrasonic processor (Model W-225, manufactured by Heat System-Ultrasonics, Inc.) for 5 minutes (30W) to obtain a substrate.

(2) Enzyme

Porcine pancreatic phospholipase A₂ [(161454.122416) manufactured by Boehringer Mannheim. Yamanouchi K.K.] was used.

(3) Measurement of phospholipase A₂ activity

To a 96 well microtitration plate (flat bottom, manufactured by Sumitomo Bakelite Medical Co., Ltd.), 40 µl of the substrate, 5 µl of a solution prepared by dissolving 10 mg of a test compound in 500 µl of dimethylsulfoxide, followed by an addition of 500 µl of a 50 mM Tris-HCl buffer solution, and 5 µl of an enzyme solution of 20 ng/ml (prepared by diluting the enzyme in a 50 mM Tris-HCl buffer solution), were added and reacted at 37° C. for 30 minutes. After termination of the reaction, the released free fatty acid was quantitatively analyzed in accordance with the ACS-ACOD (acyl CoA synthetase-acyl CoA oxidase) method [a kit of NEFA C test wako (manufactured by Wako Pure Chemical Industries, Ltd.) was used]. The quantitative analysis was made by means of Microplate ELISA Reader (Model 2550EIA Reader, manufactured by Bio-Rad Laboratories) at a wavelength of 540 nm. Separately, such experiments as mentioned above, were carried out at various concentrations (2 µg/ml, 1 µg/ml and 0.5 µg/ml) of phospholipase A₂ without a test compound. Then, the concentration of the free fatty acid versus the concentration of phospholipase A₂ was plotted.

From this standard curve, the apparent concentration of phospholipase A₂ in the case with a test compound, was read. Then, the percent inhibition of the enzyme was calculated by the following formula. The results are shown in Table 6.

$$\text{Percent inhibition (\%)} = \left(1 - \frac{A}{B}\right) \times 100$$

A: Apparent enzyme concentration when a test compound is added.

B: True enzyme concentration when a test compound is added.

TABLE 6

Compound No.	% inhibition of PLA ₂ (1,000 ppm)
1	45
2	55
3	67
4	74
5	39
8	81
9	71
10	60
11	52
12	89
13	87
14	54
15	62
16	43
17	46
18	64
19	>90
20	74
21	62
22	74
23	37
24	66
26	35
27	62
28	71

TABLE 6-continued

Compound No.	% inhibition of PLA ₂ (1,000 ppm)
29	47
30	87
32	50
38	35
39	41
41	89
43	47
44	43
45	50
46	47
47	75
48	48
49	30
50	78
51	63
52	49
53	37
54	37
55	49
57	57
58	74

TEST EXAMPLE 2

Phospholipase A₂ inhibitory activity, method B

(1) Preparation of substrate

To a solution prepared by dissolving 9.2 mg of L-α-dipalmitoylphosphatidylcholine (manufactured by Nichiyu Liposome K.K.) in 0.5 ml of chloroform, a solution prepared by dissolving 32 mg of sodium cholate (manufactured by Wako Pure Chemical Industries, Ltd.) in 0.5 ml of methanol, was added, followed by mixing. The solvent of the mixture was removed under a nitrogen stream, and then 2.5 ml of a 250 mM sodium chloride solution [prepared by dissolving sodium chloride in a 100 mM Tris-HCl buffer solution {tris(hydroxymethyl)aminomethane (manufactured by Nacalai Tesque K.K.) was adjusted to pH8.0 with hydrochloric acid}] was added thereto, and the mixture was dissolved under stirring to obtain a substrate.

(2) Enzyme

Porcine pancreatic phospholipase A₂ [(161454.122416) manufactured by Boehringer Mannheim. Yamanouchi K.K.] was used.

(3) Measurement of phospholipase A₂ activity

To a 96 well microtitration plate, 20 µl of a solution containing calcium chloride, bovine serum albumin (manufactured by Sigma Chemical, Co.) and a Tris-HCl buffer solution (pH8.0) at concentrations of 25 mM, 4.5 mg/ml and 100 mM, respectively, 5 µl of a solution prepared by dissolving 10 mg of a test compound in 500 µl of dimethylsulfoxide, followed by an addition of 500 µl of a 200 mM Tris-HCl buffer solution, 5 µl of an enzyme solution (10 µg/ml) [prepared by dissolving the enzyme in a bovine serum albumin solution (prepared by dissolving bovine serum albumin in a 100 mM Tris-HCl buffer solution at a concentration of 1 mg/ml)] and 20 µl of the substrate, were added and reacted at 37° C. for 30 minutes. After termination of the reaction, the released free fatty acid was quantitatively analyzed in accordance with the ACS-ACOD (acyl CoA synthetase-acyl CoA oxidase) method [a kit of NEFA C test wako (manufactured by Wako Pure Chemical Industries, Ltd.) was used]. The quantitative analysis was

made by means of Microplate ELISA Reader (Model 2550EIA Reader, manufactured by Bio-Rad Laboratories) at a wavelength of 540 nm. Separately, such experiments as mentioned above, were carried out at various concentrations (1 $\mu\text{g/ml}$, 0.75 $\mu\text{g/ml}$, 0.5 $\mu\text{g/mol}$ and 0.25 $\mu\text{g/ml}$) of phospholipase A_2 without a test compound. Then, the concentration of the free fatty acid versus the concentration of phospholipase A_2 was plotted.

From this standard curve, the apparent concentration of phospholipase A_2 in the case with a test compound, was read. Then, the percent inhibition of the enzyme was calculated by the following formula. The results are shown in Table 7.

$$\text{Percent inhibition (\%)} = \left(1 - \frac{A}{B}\right) \times 100$$

A: Apparent enzyme concentration when a test compound is added.

B: True enzyme concentration when a test compound is added.

TABLE 7

Compound No.	% inhibition of PLA ₂ (1,000 ppm)
7	50
10	51
13	51
18	49
19	75
43	49
44	64
45	41
47	90
53	100
58	42
60	41
61	36
62	53
63	34
64	61
65	71
66	52
67	82
68	81
69	63
70	40
71	77
72	73
73	53
74	33
75	81
76	61
77	61
78	51
79	65
80	73
81	94
82	38
83	64
84	56
85	33
86	93
87	88
88	83
89	51
90	79
91	81
92	75
93	48
94	63
95	85
97	88
98	65
99	86

TABLE 7-continued

Compound No.	% inhibition of PLA ₂ (1,000 ppm)
100	83
103	86
104	61
106	78
108	61
109	67
110	58
111	41
112	79
113	35
114	53
115	52
116	69
117	65
118	84
121	90
122	56
123	86
124	78
125	86
126	84
127	89
251	85
259	61
260	53

TEST EXAMPLE 3

Inhibitory activity on increased vascular permeability induced by acetic acid (Mouse Whittle method, method C)

Using ddy male mice, each test group consisted of 4 or 5 mice. A test compound was mixed with Tween 80 [polyoxyethylenesorbitan monooleate (manufactured by Nacalai Tesque K.K.)], and distilled water was added thereto to obtain a 2% Tween 80 suspension, or it was dissolved in the form of a salt in water to obtain an aqueous solution. A test compound was orally administered, and upon expiration of one hour from the administration, 0.7% acetic acid was intraperitoneally injected to each mouse in an amount of 0.1 ml/10 g, and at the same time, 2% brilliant blue was intravenously injected into the tail vein in an amount of 0.1 ml/20 g. Thirty minutes after the injection of brilliant blue, the cervical vertebrae were dislocated under anesthesia by chloroform, and the abdominal cavity was washed with 5 ml of a physiological saline. The washing solution was subjected to centrifugal separation at 3,000 rpm for 10 minutes, and the amount of the dye in the supernatant was measured at 600 nm absorbance by Microplate ELISA Reader (Model 2550EIA Reader, manufactured by Bio-Rad Laboratories). The inhibition rate of the amount of leaked dye in the group in which a test compound was administered relative to the control group was obtained by the following formula. The results are shown in Table 8.

$$\text{Inhibition rate (\%)} = \left(1 - \frac{C}{D}\right) \times 100$$

C: Amount of leaked dye in the group to which a test compound was administered.

D: Amount of leaked dye in the control group.

TABLE 8

Compound No.	Dose (mg/kg)	Inhibition rate (%)
1	50	46
2	20	51
3	50	58
4	50	43
5	50	53
7	20	53
8	20	48
9	50	81
10	25	53
	10	42
11	100	49
13	100	57
15	50	41
16	20	55
17	50	31
18	25	49
20	20	48
22	20	81
	10	39
23	20	33
39	20	53
41	100	85
43	20	48
45	20	29
47	20	72
	10	46
49	20	50
55	25	59
57	20	43
63	10	41
78	20	51
	10	32
79	20	67
86	20	42
87	10	28
93	20	47
	10	40
94	20	53
101	20	46
120	20	43
251	20	43

TEST EXAMPLE 4

Inhibitory-activity on increased vascular permeability induced by acetic acid (Rat Whittle method, method D

Using SD (Crj: CD) male rats, each test group consisted of from 3 to 5 rats. A test compound was mixed with Tween 80 [polyoxyethylenesorbitan monooleate (manufactured by Nacalai Tesque K.K.)], and distilled water was added thereto to obtain a 2% Tween 80 suspension, or it was dissolved in the form of a salt in water to obtain an aqueous solution. A test compound was orally administered, and one hour later, 0.7% acetic acid was intraperitoneally injected to each rat in an amount of 0.05 ml/10 g, and at the same time, 2% brilliant blue was intravenously injected into the tail vein in an amount of 0.05 ml/20 g. Thirty minutes after the injection of brilliant blue, the cervical vertebrae were dislocated under anesthesia by chloroform, and the abdominal cavity was washed with 10 ml of a physiological saline. The washing solution was subjected to centrifugal separation at 3,000 rpm for 10 minutes, and the amount of the dye in the supernatant was measured at 600 nm absorbance by Microplate ELISA Reader (Model 2550EIA Reader, manufactured by Bio-Rad Laboratories). The inhibition rate of the amount of leaked dye in the group to which a test compound was administered relative to the control group was obtained

from the following formula, and the results are shown in Table 9.

$$\text{Inhibition rate (\%)} = \left(1 - \frac{C}{D}\right) \times 100$$

C: Amount of leaked dye in the group to which a test compound was administered.

D: Amount of leaked dye in the control group.

TABLE 9

Compound No.	Dose (mg/kg)	Inhibition rate (%)
2	100	38
3	100	75
10	100	57
	50	37
16	100	96
17	50	40
19	50	34
20	100	49
22	100	58
23	100	40
43	50	72
45	50	27
46	50	31
47	50	82
	25	56
49	50	30
55	25	69
	12.5	43
57	50	47
58	50	31
60	50	72
61	25	61
63	50	39
	25	31
66	25	72
69	25	48
72	25	66
78	50	55
	25	40
79	50	74
80	50	35
	25	33
82	25	38
86	50	37
87	25	61
	12.5	47
93	50	71
	25	54
94	50	55
	25	45
98	50	32
101	50	41
113	50	67
120	100	56
	50	35
121	12.5	31
251	25	70
	12.5	46

TEST EXAMPLE 5

Inhibitory activity on carrageenin edema

Using Wister male rats (body weight: about 100 g), each test group consisted of 5 rats. A test compound was mixed with Tween 80 [polyoxyethylenesorbitan monooleate (manufactured by Nacalai Tesque K.K.)], and distilled water was added thereto to obtain a 2% Tween 80 suspension, or it was dissolved in the form of a salt in water to obtain an aqueous solution. Either the suspension or the aqueous solution was orally administered in an amount of 200 mg/kg, 100 mg/kg, 50 mg/kg or 25 mg/kg. One hour later, 0.1 ml of a 1% λ-carra-

geenin solution dissolved in a physiological saline was injected subcutaneously to the right hind paw of each rat to cause inflammation. Three hours later, the paw volume was measured by a paw volume measuring device (manufactured by Ugobasiee K.K.). A swelling volume was obtained from the difference from the value before the inflammation. The inhibition rate was calculated by the following formula, and the results are shown in Table 10.

$$\text{Inhibition rate (\%)} = \left(1 - \frac{F}{E}\right) \times 100$$

F: Average swelling volume in the group to which a test compound was administered.

E: Average swelling volume in the control group.

TABLE 10

Compound No.	Dose (mg/kg)	Inhibition rate (%)	
2	100	17	
3	100	20	
5	100	37	
10	100	28	
11	100	24	
13	100	21	
16	100	24	
19	100	31	
22	100	29	
23	100	30	
25	200	27	
28	50	25	
39	100	25	
43	50	31	
45	50	23	
46	50	30	
47	50	41	
57	100	35	
60	50	27	
65	50	37	
66	50	31	
67	25	19	
69	50	25	
72	25	21	
73	50	20	
77	50	22	
78	50	26	
79	50	20	
80	50	29	
82	50	19	
86	50	27	
87	50	21	
91	50	23	
93	50	22	
94	50	23	
98	50	45	
101	50	24	
104	50	48	
106	50	19	
110	50	25	
113	50	26	
114	50	28	
120	50	27	
123	50	42	
125	50	22	
150	50	23	
251	50	30	
259	50	17	

TEST EXAMPLE 6

Acute toxicity

Administration route: Intravenous injection

Using ddy male mice (body weight: 25–30 g), each test group consisted of 5 mice. A test compound was

dissolved in the form of a sodium salt in a physiological saline or in a 5% glucose aqueous solution, and intravenously injected in an amount of 0.1 ml/10 g body weight. After the injection, the mortality rate was obtained over one week, and the median lethal dose LD₅₀ (mg/kg) was determined. The results are shown in Table 11.

TABLE 11

Compound No.	LD ₅₀ (mg/kg)	
1	100~150	
2	50~100	
3	>100	
8	>25	
9	>150	
10	50~100	
11	>150	
12	>150	
13	>70	
15	100~150	
16	>100	
17	50~100	
18	>150	
19	50~100	
21	>75	
22	>100	
24	>150	
40	50~100	
43	78	
45	98	
47	58	
49	175	
55	237	
57	83	
60	>60	
61	>80	
63	>130	
68	>80	
73	>80	
77	>80	
78	>60	
80	>80	
86	>40	
87	75	
91	>80	
106	>20	
120	83	
251	65	

TEST EXAMPLE 7

Effects against acute pancreatitis

Using Crj-CD male rats (for Compound No. 19, rats having a body weight of from 371 to 484 g were used, and for Compound No. 10, rats having a body weight of from 444 to 574 g were used), each test group consisted of 3 rats. An experimental acute pancreatitis model was prepared by a closed duodenal loop method under anesthesia with halothane (manufactured by Hoechst Japan) and nitrous oxide (manufactured by Sumitomo Seika K.K.) applied by means of a general inhalation anesthesia machine (Model EM-2 and halothane evaporator F-Model). Then, Compound No. 19 or Compound No. 10 was continuously intravenously injected into the tail vein in an amount of 50 mg per kg or 40 mg per kg, respectively, at a rate of 0.05 ml per minute by means of a pump (Technicon AA II Proportioning Pump III, manufactured by Technicon Instruments Corporation). No injection was made to a control group. Gross pathological examination was conducted upon expiration of 6 hours after the ventrotomy in the case of the test group to which Compound No. 19 was administered, or upon expiration of 3 hours after the ventrotomy in the case of

the test group to which Compound No. 10 was administered. As a result, as shown in the following Table 12, the groups to which the compounds of the present invention were administered, show distinct usefulness for treating acute pancreatitis.

TABLE 12

Groups	Pancreatic hemorrhage Petechia		Pancreatic edema	
	Grade	Distribution	Grade	Distribution
Control group	++	++	++	++
(against the group to which Compound No. 19 was administered)	++	++	+++	++
Group to which Compound No. 19 was administered	+++	+++	+++	++
Control group	-	-	+	+
(against the group to which Compound No. 10 was administered)	-	-	++	++
Group to which Compound No. 10 was administered	++	++	++	++
Control group	+	+	++	++
(against the group to which Compound No. 10 was administered)	±	±	++	++
Group to which Compound No. 10 was administered	±	±	±	±
Control group	+	+	+	+
(against the group to which Compound No. 10 was administered)	+	+	++	+

Grade of pancreatic lesions

-: No significant lesions,

±: Minimal,

+: Light,

++: Moderate,

+++ : Marked

Distribution of pancreatic lesions

-: No significant lesions,

± ~ +++ : Focal-diffuse

TEST EXAMPLE 8

Effects against acute pancreatitis

Using Crj-CD male rats, each test group consisted of 3 rats. An experimental acute pancreatitis model was prepared by a closed duodenal loop method under anesthesia with halothane (manufactured by Hoechst Japan) and nitrous oxide (manufactured by Sumitomo Seika K.K.) applied by a general inhalation anesthesia machine (Model EM2 and halothane evaporator F-Model). Each compound (subjected to the test in the form of a sodium salt) was continuously intravenously injected into the tail vein in an amount of 0.4 ml/100 g to 0.6 ml/100 g at a rate of 0.05 ml per minute by a pump (Technicon AA II Proportioning Pump III, manufactured by Technicon Instruments Corporation) or rapidly intravenously injected. No injection was made to a control group. Gross pathological examination was conducted upon expiration of 6 hours after the ventrotomy in the case of the group to which the compound was administered. With respect to each of four lesions among pancreatic lesions i.e. petechia, ecchymosis, pancreatic necrosis and abdominal fatty necrosis, the grade and the distribution of lesions were scored with five grades of 0, 0.5, 1, 2 and 3 (severe lesions are 3). The sum of all lesions was designated as scores of pancreatitis lesions, and the sum of the score of petechia and the score of ecchymosis only was designated as scores of hemorrhagic lesions. The pancreatitis inhibition rate (%) and the hemorrhage inhibition rate (%) were obtained by the following formulas, and the results are shown in Table 13.

$$\text{Pancreatitis inhibition rate (\%)} = \left(1 - \frac{H}{G}\right) \times 100$$

H: Scores of pancreatitis lesions of the group to which a test compound was administered.

G: Scores of pancreatitis lesions of the control group.

$$\text{Hemorrhage inhibition rate (\%)} = \left(1 - \frac{J}{I}\right) \times 100$$

J: Scores of hemorrhagic lesions of the group to which a test compound was administered.

I: Scores of hemorrhagic lesions of the control group.

TABLE 13

Compound No.	Dose (mg/kg)	*1	*2
1	10	66	49
2	26*	46	
3	10	49	51
9	10	36	21
11	23*	52	
13	23*	100	
14	19*	52	
15	10	45	61
16	20*	52	
17	20*	73	
21	27*	57	
24	11*	68	
34	10	30	30
35	10	35	35
43	20*	81	
45	25*	62	
46	46*	36	
47	20*	68	
49	42*	68	
55	40*	65	
57	20*	60	
58	10	70	51
60	10	92	94
61	10	79	64
62	10	45	61
63	10	83	66
64	10	60	68
65	10	67	74
66	10	53	63
68	10	74	77
72	10	62	32
73	10	74	79
74	10	66	67
77	10	66	70
78	10	96	91
79	10	23	39
80	10	11	8
81	10	49	58
83	10	53	51
85	10	57	67
86	10	87	85
87	10	83	87
93	10	70	70
94	10	11	11
97	10	35	35
106	10	96	97
107	10	63	61
113	10	41	36
114	10	32	27
117	10	30	30
120	24*	100	
122	10	51	51
123	10	56	56
124	10	51	51
251	10	79	80

Note:

Symbol * in the column for "Dose" indicates a case of continuous intravenous injection, and no symbol indicates a case of single intravenous injection.

*1: Inhibition rate of hemorrhagic lesions (%)

*2: Inhibition rate of pancreatitis lesions (%)

To administer the compound of the present invention for the treatment of the above-mentioned diseases

caused by phospholipase A₂, it is formulated alone or together with a pharmaceutically acceptable carrier into a drug composition suitable for peroral, or parenteral administration, such as a tablet, a powder, a capsule, a granule, an injection drug, an ointment, an inhalant or a suppository, and it is administered in the form of such a drug formulation.

As a drug formulation suitable for peroral administration, a solid composition such as a tablet, a capsule, a powder, a granule or a troach; or a liquid composition such as a syrup suspension, may be mentioned. The solid composition such as a tablet, a capsule, a powder, a granule or a troach may contain a binder such as fine crystalline cellulose, gum arabic, tragacanth gum, gelatine or polyvinyl chloride; an excipient such as starch, lactose or carboxymethyl cellulose; a disintegrator such as arginic acid, corn starch or carboxymethyl cellulose; a lubricant such as magnesium stearate, light silicic anhydride or colloidal silicon dioxide; a sweetener such as sucrose; or a flavoring agent such as peppermint or methyl salicylate. The liquid composition such as a syrup or a suspension may contain sorbitol, gelatine, methyl cellulose, carboxymethyl cellulose, a vegetable oil such as a peanut oil, an emulsifier such as lecithin as well as a sweetener, a preservative, a colorant or a flavoring agent, as the case requires. Such a composition may be provided in the form of a dried formulation. These formulations preferably contain from 1 to 95% by weight of the active compound.

A drug formulation suitable for parenteral administration may, for example, be an injection drug. The injection drug may be prepared by dissolving the compound in the form of a salt in usual water for injection, or may be formulated into a formulation suitable for injection such as a suspension or an emulsion (in a mixture with a pharmaceutically acceptable oil or liquid). In such a case, it may contain benzyl alcohol as an antibacterial agent, ascorbic acid as an antioxidant, a pharmaceutically acceptable buffer solution or a reagent for adjusting the osmotic pressure. Such an injection drug preferably contains from 0.1 to 8% by weight of the active compound.

A drug formulation suitable for topical or per rectal administration may, for example, be an inhalant, an ointment or a suppository. The inhalant may be formulated by dissolving the compound of the present invention alone or together with a pharmaceutically acceptable inert carrier in an aerosol or nebulizer solution, or may be administered to the respiratory airway in the form of fine powder for inhalation. In the case of fine powder for inhalation, the particle size is usually not more than 50 μ m, preferably not more than 10 μ m. Such an inhalant may be used, if necessary, in combination with other antiasthmatic agent or bronchodilator.

An ointment may be prepared by a conventional method by an addition of a commonly employed base or the like. The ointment preferably contains from 0.1 to 30% by weight of the active compound.

The suppository may contain a carrier for formulation which is well known in this field, such as polyethylene glycol, lanolin, cacao butter or fatty acid triglyceride. The suppository preferably contains from 1 to 95% by weight of the active compound.

The above-mentioned drug compositions suitable for peroral, parenteral, topical or per rectal administration, may be formulated by conventional methods so that after administration to a patient, the active component

will be rapidly discharged, gradually discharged or belatedly discharged.

The dose of the compound of the present invention varies depending upon the type of the compound, the administration method, the condition of the patient or the animal to be treated. The optimum dose and the number of administration under a specific condition must be determined by the judgement of a competent doctor. Usually, however, a daily dose to an adult is from about 0.01 g to about 10 g, preferably from about 0.05 g to about 5 g. In the case of the above inhalation method, the dose of the compound of the present invention is preferably from about 0.01 mg to about 100 mg per administration.

Now, specific Formulation Examples of the phospholipase A₂ inhibitor, the anti-inflammatory agent or the anti-pancreatitis agent of the present invention will be given.

FORMULATION EXAMPLE 1 (tablet)

(1) Compound No. 30	200 mg
(2) Lactose	150 mg
(3) Starch	30 mg
(4) Magnesium stearate	6 mg

The above composition is tableted so that the components (1) to (4) constitute one tablet.

FORMULATION EXAMPLE 2 (powder or microgranule)

(1) Compound No. 35	200 mg
(2) Sugar ester (DK ester F-160, manufactured by Daiichi Kogyo)	180 mg
(3) Surfactant (Dekagreen I-L, manufactured by Nikko Chemicals)	15 mg
(4) Light silicic anhydride	25 mg

The component (1) is wet-pulverized in an aqueous solution containing 5% of the component (3). Then, 180 mg of the component (2) is added thereto, and the mixture is freeze-dried. The dried product is pulverized and mixed with the component (4).

The mixture is formed into a power or microgranule. Such a powder or microgranule may be sealed in a capsule to obtain a capsule drug.

FORMULATION EXAMPLE 3 (hard gelatine capsule)

(1) Sodium salt of Compound No. 10	250 mg
(2) Starch	200 mg
(3) Magnesium stearate	10 mg

The components (1) to (3) is packed in a hard gelatine capsule to obtain a hard gelatine capsule drug.

FORMULATION EXAMPLE 4 (injection drug)

(1) Sodium salt of Compound No. 19	1 g
(2) Glucose	10 g
(3) Distilled water for injection	200 ml

The components (1) to (3) are formulated into an injection drug in accordance with a usual method for preparation of an injection drug.


FORMULATION EXAMPLE 5
(ointment for external skin application)

(1) Sodium salt of Compound No. 10	5 g
(2) White vaseline	25 g
(3) Stearyl alcohol	22 g
(4) Propylene glycol	12 g
(5) Sodium lauryl sulfate	1.5 g
(6) Ethyl para-hydroxybenzoate	0.025 g
(7) Propyl para-hydroxybenzoate	0.015 g
(8) Purified water	100 g

The components (1) to (8) are formulated into an ointment for external skin application by a usual method for preparation of an ointment.

We claim:

1. A diaminotrifluoromethylpyridine derivative of the formula (I) or its salt:



wherein X is $\text{—CW}^1\text{R}^1$ and Y is $\text{—(NH)}_m\text{SO}_2\text{R}^9$; wherein W is an oxygen or sulfur atom, m is 0 or 1, and R^1 and R^9 are the same or different and are alkyl, alkenyl, alkenyl, alkenyl, cycloalkyl, cycloalkenyl, phenyl, or a member selected from the group consisting of naphthyl, tetrahydronaphthyl, indanyl, adamantyl, noradamantyl, norbornanyl and norbornanonyl; wherein each of R^1 and R^9 is optionally substituted with a member selected from the group consisting of a halogen, alkyl, haloalkyl, alkoxy, haloalkoxy, alkylthio, cycloalkyl, cycloalkoxy, cycloalkenyl, cycloalkenyloxy, alkoxycarbonyl, alkylcarbonyl, alkylcarbonyloxy, aryl, aryloxy, arylthio, amino, alkylsub-

stituted amino, alkylsubstituted cyano and alkylsubstituted nitro.

2. The diaminotrifluoromethylpyridine derivative or its salt according to claim 1, wherein R¹ is alkyl, haloalkyl, alkenyl, haloalkenyl, cycloalkyl, halogen-substituted cycloalkyl, phenyl, halogen-substituted phenyl, alkyl- or haloalkyl-substituted phenyl, or alkoxy- or haloalkoxy-substituted phenyl, and R² is alkyl, haloalkyl, phenyl, halogen-substituted phenyl, alkyl- or haloalkyl-substituted phenyl or alkoxy- or haloalkoxy-substituted phenyl.

3. The diaminotrifluoromethylpyridine derivative or its salt according to claim 1, wherein the diaminotrifluoromethylpyridine derivative is at least one derivative selected from the group consisting of N-(2-ethylsulfonylamino-5-trifluoromethyl-3-pyridyl)cyclohexanecarboxamide, N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)-5-indanecarboxamide, N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)-acetoxycetamide, N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)crotonamide, N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)-3-trifluoromethylbenzamide, N-(2-ethylsulfonylamino-5-trifluoromethyl-3-fluorobenzamide, N-(2-methylsulfonylamino-5-trifluoromethyl-3-pyridyl)-6-(1,2,3,4-tetrahydronaphthalene)carboxamide and N-(2-ethylsulfonylamino-5-trifluoromethyl-3-pyridyl)crotonamide.

4. The diaminotrifluoromethylpyridine derivative or its salt according to claim 1, wherein the diaminotrifluoromethylpyridine derivative is N-(2-ethylsulfonylamino-5-trifluoromethyl-3-pyridyl)cyclohexanecarboxamide.

5. A pharmaceutical composition for use as a phospholipase A₂ inhibitor, an anti-inflammatory agent or an anti-pancreatic agent which contains as an active ingredient said diaminotrifluoromethylpyridine derivative according to claim 1 and a carrier.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 5,229,403
DATED : July 20, 1993
INVENTOR(S) : Takahiro Haga et al.

It is certified that error appears in the above-indentified patent and that said Letters Patent is hereby corrected as shown below:

On the title page, Item [30],

The second Foreign Application Priority Data has been omitted,
please insert: --May 24, 1991 [JP] Japan.....3-222530--

Signed and Sealed this
Twenty-second Day of March, 1994

Attest:



BRUCE LEHMAN

Attesting Officer

Commissioner of Patents and Trademarks

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 5,229,403
DATED : July 20, 1993
INVENTOR(S) : Takahiro Haga et al.

It is certified that error appears in the above-indentified patent and that said Letters Patent is hereby corrected as shown below:

On the title page, Item [30],

The second Foreign Application Priority Data has been omitted,
please insert: --May 24, 1991 [JP] Japan.....3-222530--

Signed and Sealed this
Twenty-second Day of March, 1994

Attest:



BRUCE LEHMAN

Attesting Officer

Commissioner of Patents and Trademarks